


# Chapter Opening Features

## Chapter Opening Photos

Each chapter begins with photos and captions that are related to an Excursion in the chapter.

## Web Icon

The web icon on this opening page lets students know of additional online resources at [math.college.hmco.com/students](http://math.college.hmco.com/students).



page 322

The Tower of Hanoi is a puzzle that has the following form. Three pegs are placed on a board. A number of disks, graded in size, are stacked on one of the pegs with the largest disk on the bottom and successively smaller disks placed on top. The disks are moved according to the following rules:

1. Only one disk at a time may be moved.
2. A larger disk may not be placed on top of a smaller disk.

The object of the game is to transfer all of the disks, one at a time, from one peg to one of the other two pegs. If initially there is only one disk, then only one move is required. With two disks, three moves are required; with three disks, seven moves are required.

The chart below shows the minimum number of moves required for a given number of disks. The increase in the number of moves required for each additional disk is also given.

Number of disks	1	2	3	4	5	6	7
Minimum number of moves	1	3	7	15	31	63	127
Increase in number of moves		$2 \cdot 1 = 2$	$2 \cdot 3 = 4$	$2 \cdot 7 = 8$	$2 \cdot 15 = 18$	$2 \cdot 31 = 32$	$2 \cdot 63 = 64$

**point of interest**  
If recent estimates of the age of the universe are accurate, our solar system is about 4.5 billion years old. The first known life forms occurred about 3 billion years ago.

For the list of numbers in the bottom row of the table, each successive number can be found by multiplying the preceding number by a constant (in this case 2). This list of numbers can be represented by the equation  $f(n) = 2^n$ , which is an example of an exponential function, one of the topics of this chapter.

The formula for the minimum number of moves is given by  $M = 2^n - 1$ , which contains the exponential expression  $2^n$ . In this formula,  $M$  is the minimum number of moves required to transfer  $n$  disks to one of the other pegs.

There is an ancient myth involving the Tower of Hanoi puzzle and the lifetime of the universe. In this myth, three priests sit in the center of the universe with 3 diamond needles and 64 golden disks on one of the needles. The only job of the priests is to transfer the golden disks to one of the other needles using the rules of the Tower of Hanoi puzzle. The priests can transfer one disk to another needle every second. According to the myth, the universe will cease to exist at the precise moment the priests have completed the transfer of all 64 disks to one of the other needles.

Thus according to the myth, the lifetime of the universe is given by  $2^{64} - 1$  seconds. Use a calculator to show that this amounts to approximately 585 billion years! Even if the priests started the transfer of the disks 12 billion years ago (when astronomers estimate our universe began), the myth indicates that the universe will continue to exist for another 573 billion years.

323

## Chapter Opener Subject Matter

The second page of each chapter opener presents a motivational topic, an application from the chapter, or a new mathematical concept.

page 323





## Excursion

## Earned Run Average

One measure of a pitcher's success is earned run average. **Earned run average (ERA)** is the number of earned runs a pitcher gives up for every nine innings pitched. The definition of an earned run is somewhat complicated, but basically an earned run is a run that is scored as a result of hits and base running that involves no errors on the part of the pitcher's team, if the opposing team scores a run on an error (for example, a fly ball that should have been caught in the outfield was fumbled), then that run is not an earned run.

A proportion is used to calculate a pitcher's ERA. Remember that the statistic involves the number of earned runs per nine innings. The answer is always rounded to the nearest hundredth. Here is an example.

During the 2001 regular baseball season, Chan Ho Park gave up 91 earned runs and pitched 234 innings for the Los Angeles Dodgers. To calculate Chan Ho Park's ERA, let  $x$  = the number of earned runs for every nine innings pitched. Write a proportion and then solve for  $x$ .

$$\frac{91 \text{ earned runs}}{234 \text{ innings}} = \frac{x \text{ earned runs}}{9 \text{ innings}}$$

$$91 \cdot 9 = 234 \cdot x$$

$$819 = 234x$$

$$\frac{819}{234} = \frac{234x}{234}$$

$$3.5 = x$$

Chan Ho Park's ERA was 3.50.

## Excursion Exercises

- In 1979, his rookie year, Jeff Reardon pitched 21 innings for the New York Mets and gave up four earned runs. Calculate Reardon's ERA for 1979.
- Roger Clemens's first year with the Boston Red Sox was 1984. During that season, he pitched 133.1 innings and gave up 64 earned runs. Calculate Clemens's ERA for 1984.
- During the 1998 baseball season, Pedro Martinez of the Boston Red Sox pitched 233.2 innings and gave up 75 earned runs. During the 1999 season, he gave up 49 earned runs and pitched 213.1 innings. During which season was his ERA lower? How much lower?
- In 1987, Nolan Ryan had the lowest ERA of any pitcher in the major leagues. He gave up 65 earned runs and pitched 211.2 innings for the Houston Astros. Calculate Ryan's ERA for 1987.
- Find the necessary statistics for a pitcher on your "home team," and calculate that pitcher's ERA.

Earned Run Average Leaders		
Year	Player, club	ERA
National League		
1990	Dwight Gooden, Houston	2.21
1991	Dennis Martinez, Montreal	2.39
1992	Bill Swift, San Francisco	2.06
1993	Greg Maddux, Atlanta	2.36
1994	Greg Maddux, Atlanta	1.56
1995	Greg Maddux, Atlanta	1.63
1996	Kevin Brown, Florida	1.89
1997	Pedro Martinez, Montreal	1.90
1998	Greg Maddux, Atlanta	2.22
1999	Randy Johnson, Arizona	2.48
2000	Kevin K. Brown, Los Angeles	2.58
2001	Randy Johnson, Arizona	2.49
American League		
1990	Roger Clemens, Boston	1.33
1991	Roger Clemens, Boston	2.62
1992	Roger Clemens, Boston	2.41
1993	Kevin Appier, Kansas City	2.56
1994	Steve Ontiveros, Oakland	2.65
1995	Randy Johnson, Seattle	2.48
1996	Juan Guzman, Toronto	2.93
1997	Roger Clemens, Toronto	2.05
1998	Roger Clemens, Toronto	2.65
1999	Pedro Martinez, Boston	2.07
2000	Pedro Martinez, Boston	1.74
2001	Freddy Garcia, Seattle	3.02

# Interactive Method, *continued*

## Excursions

Each section ends with an Excursion along with corresponding Excursion Exercises. These activities engage students in the mathematics of the section. Some Excursions are designed as in-class cooperative learning activities that lend themselves to a hands-on-approach. They can also be assigned as projects or extra credit assignments. The Excursions provide opportunities for student to take an active role in the learning process. The photos on the first page of a chapter opener relate to one of the Excursions in that chapter.

## AIM for Success Student Preface

This 'how to use this text' preface explains what is required of a student to be successful and how this text has been designed to foster student success.

AIM for Success can be used as a lesson on the first day of class or as a project for students to complete to strengthen their study skills.



Charles Dodgson (Lewis Carroll)

**Math Matters** Charles Dodgson

One of the most well known logicians is Charles Dodgson (1832–1898). Dodgson was educated at Rugby and Oxford, and in 1861 he became a lecturer in mathematics at Oxford. Some of his mathematical works include *A Syllabus of Plane Algebraical Geometry*, *The Fifth Book of Euclid Treated Algebraically*, and *Symbolic Logic*. Although Dodgson was a distinguished mathematician in his time, he is best known by his pen name Lewis Carroll, which he used when he published *Alice's Adventures in Wonderland* and *Through the Looking Glass*.

Queen Victoria of the United Kingdom enjoyed *Alice's Adventures in Wonderland* to the extent that she told Dodgson she was looking forward to reading another of his books. He promptly sent her his *Syllabus of Plane Algebraical Geometry*; and it was reported that she was less than enthusiastic about the latter book.



**Compound Statements**

Connecting statements with words and phrases such as *and*, *or*, *not*, *if ... then*, and *if and only if* creates a **compound statement**. For instance, "I will attend the meeting or I will go to school" is a compound statement. It is composed of the two **component statements** "I will attend the meeting" and "I will go to school." The word *or* is a **connective** for the two component statements.

George Boole used symbols such as *p*, *q*, *r*, and *s* to represent statements and the symbols  $\wedge$ ,  $\vee$ ,  $\neg$ ,  $\rightarrow$ ,  $\leftrightarrow$ , and  $\leftrightarrow$  to represent connectives. See Table 3.1.

Table 3.1 Logic Symbols

Original Statement	Connective	Statement in Symbolic Form	Type of Compound Statement
not <i>p</i>	not	$\neg p$	negation
<i>p</i> and <i>q</i>	and	$p \wedge q$	conjunction
<i>p</i> or <i>q</i>	or	$p \vee q$	disjunction
if <i>p</i> , then <i>q</i>	if ... then	$p \rightarrow q$	conditional
<i>p</i> if and only if <i>q</i>	if and only if	$p \leftrightarrow q$	biconditional

**QUESTION** What connective is used in a conjunction?

**ANSWER** The connective *and*.

# Math Matters and Margin Notes

## Math Matters

This feature typically contains an interesting sidelight about mathematics, its history, or its applications.

## Historical Note

These margin notes provide historical background information related to the concept under discussion or vignettes of individuals who were responsible for major advancements in their fields of expertise.

## Calculator Note

These notes provide information about how to use the various features of a calculator.

page 112

**The Natural Exponential Function**

For all real numbers *x*, the function defined by  $f(x) = e^x$  is called the **natural exponential function**.

A calculator with an  $e^x$  key can be used to evaluate  $e^x$  for specific values of *x*. For instance,

$$e^2 \approx 7.389056, \quad e^{3.3} \approx 33.115452, \quad \text{and} \quad e^{-1.4} \approx 0.246597$$

The graph of the natural exponential function can be constructed by plotting a few points or by using a graphing utility.

**EXAMPLE 3** Graph a Natural Exponential Function


Graph  $f(x) = e^x$ .

**Solution**


Use a calculator to find range values for a few domain values. The range values in the table below have been rounded to the nearest tenth.

<i>x</i>	-2	-1	0	1	2
$f(x) = e^x$	0.1	0.4	1.0	2.7	7.4

Plot the points given in the table and then connect the points with a smooth curve. Because  $e > 1$ , as *x* increases,  $e^x$  increases. Thus the values of *y* increase as *x* increases. As *x* decreases,  $e^x$  becomes closer to zero. For instance, when  $x = -5$ ,  $e^{-5} \approx 0.0067$ . Thus as *x* decreases, the graph gets closer and closer to the *x*-axis. The *y*-intercept is (0, 1).



In the figure at the right, compare the graph of  $f(x) = e^x$  with the graphs of  $g(x) = 2^x$  and  $h(x) = 3^x$ . Because  $2 < e < 3$ , the graph of  $f(x) = e^x$  is between the graphs of *g* and *h*.



**CHECK YOUR PROGRESS 3** Graph  $f(x) = e^{-x} + 2$ .

**Solution** See page S23.

**historical note**



**Leonhard Euler**

(1707–1783)


Some mathematicians consider Euler to be the

greatest mathe-

matician of all time. He certainly was the most prolific writer of mathematics of all time. He was the first to introduce many of the mathematical notations that we use today. For instance, he introduced the symbol  $\pi$  for pi, the functional notation  $f(x)$ , and the letter *e* as the base of the natural exponential function. ■

**CALCULATOR NOTE**

The graph below was produced on a TI-83 graphing calculator by entering  $e^x$  in the *Y=* menu.



# Margin Notes, *continued*

## Point of Interest

These notes provide interesting information related to the topics under discussion. Many of these are of a contemporary nature and, as such, they provide students with the needed motivation for studying concepts that may at first seem abstract and obscure without this information.

### historical note


The Babylonians knew that Earth was in approximately the same position in the sky every 365 days. Historians suggest that the reason one complete revolution of a circle is  $360^\circ$  is that 360 is the closest number to 365 that is divisible by many natural numbers. ■

### point of interest



The Leaning Tower of Pisa is the bell tower of the Cathedral in Pisa, Italy. Its construction began on August 5, 1173 and continued for about 200 years. The tower was designed to be vertical, but it started to lean during its construction. By 1350 it was 2.5° off from the vertical; by 1817, it was 5.1° off; and by 1990, it was 5.5° off. In 2001, work on the structure that returned its list to 5° was completed. (Source: Time magazine, June 25, 2001, pp. 34–35.)


In the figure at the right,  $\angle x$  and  $\angle QRS$  are two different names for the same angle.  $\angle y$  and  $\angle SRT$  are two different names for the same angle. Note that in this figure, more than two rays meet at  $R$ . In this case, the vertex alone cannot be used to name  $\angle QRT$ .



An angle is often measured in degrees. The symbol for degrees is a small raised circle,  $^\circ$ . The angle formed by rotating a ray through a complete circle has a measure of  $360^\circ$ .



A protractor is often used to measure an angle. Place the center of the protractor at the vertex of the angle with the edge of the protractor along a side of the angle. The angle shown in the figure below measures  $38^\circ$ .




A  $90^\circ$  angle is called a **right angle**. The symbol  $\perp$  represents a right angle.



**Perpendicular lines** are intersecting lines that form right angles.



The symbol  $\perp$  means "is perpendicular to." In the figure at the right,  $p \perp q$  and  $\overline{AB} \perp \overline{CD}$ . Note that line  $p$  contains  $\overline{AB}$  and line  $q$  contains  $\overline{CD}$ . Perpendicular lines contain perpendicular line segments.



This illustrates the following theorem regarding the solutions of a quadratic equation.

### The Sum and Product of the Solutions of a Quadratic Equation

If  $s_1$  and  $s_2$  are the solutions of a quadratic equation of the form  $ax^2 + bx + c = 0$ ,  $a \neq 0$ , then

$$\text{the sum of the solutions } s_1 + s_2 = -\frac{b}{a}, \text{ and}$$

$$\text{the product of the solutions } s_1 s_2 = \frac{c}{a}$$

### TAKE NOTE

The result is the same if we let  $s_1 = 6$  and  $s_2 = -2$ .

In this section, the method we used to check the solutions of a quadratic equation was to substitute the solutions back into the original equation. An alternative method is to use the sum and product of the solutions.

For example, let's check that  $-2$  and  $6$  are the solutions of the equation  $x^2 - 4x - 12 = 0$ . For this equation,  $a = 1$ ,  $b = -4$ , and  $c = -12$ . Let  $s_1 = -2$  and  $s_2 = 6$ .

$$\begin{array}{l} s_1 + s_2 = -\frac{b}{a} \\ -2 + 6 = \frac{-(-4)}{1} \\ 4 = 4 \end{array} \quad \begin{array}{l} s_1 s_2 = \frac{c}{a} \\ (-2)(6) = \frac{-12}{1} \\ -12 = -12 \end{array}$$

### The solutions check.

In Example 4, we found that the exact solutions of the equation  $2x^2 = 4x - 1$  are  $\frac{2 + \sqrt{2}}{2}$  and  $\frac{2 - \sqrt{2}}{2}$ . Use the sum and product of the solutions to check these solutions.

Write the equation in standard form. Then determine the values of  $a$ ,  $b$ , and  $c$ .

$$\begin{aligned} 2x^2 &= 4x - 1 \\ 2x^2 - 4x + 1 &= 0 \\ a &= 2, b = -4, c = 1 \end{aligned}$$

$$\text{Let } s_1 = \frac{2 + \sqrt{2}}{2} \text{ and } s_2 = \frac{2 - \sqrt{2}}{2}.$$

$$\begin{array}{l} s_1 + s_2 = -\frac{b}{a} \\ \frac{2 + \sqrt{2}}{2} + \frac{2 - \sqrt{2}}{2} = \frac{-(-4)}{2} \\ \frac{2 + \sqrt{2} + 2 - \sqrt{2}}{2} = 2 \\ 4 = 4 \end{array} \quad \begin{array}{l} s_1 s_2 = \frac{c}{a} \\ \left(\frac{2 + \sqrt{2}}{2}\right)\left(\frac{2 - \sqrt{2}}{2}\right) = \frac{1}{2} \\ \frac{4 - 2}{4} = \frac{1}{2} \\ \frac{2}{4} = \frac{1}{2} \\ \frac{1}{2} = \frac{1}{2} \end{array}$$

### The solutions check.

(continued)

If you need to review material on adding and multiplying radical expressions, see Lessons 9.2B and 9.2C on the CD that you received with this book.

## Take Note

These margin notes alert students to a point requiring special attention or are used to amplify the concepts that are currently being developed. Some Take Notes, identified by <symbol>, reference the student CD. A student who needs to review a prerequisite skill or concept can find the needed material on this CD.

# Exercises

## Exercise Sets

The exercise sets of Mathematical Excursions were carefully written to provide a wide variety of exercises that range from drill and practice to interesting challenges. Exercise sets emphasize skill building, skill maintenance, concepts, and applications, when they are appropriate. Icons are used to identify various types of exercise.

- a. Find an exponential regression equation for these data, using  $t = 0$  to represent 1995. Round to the nearest hundredth.  
 b. Use the equation to predict the number of ATMs in 2010.
37. The table below shows the saturation of water in air at various air temperatures.

Temperature (in °C)	0	5	10	20	25	30
Saturation (in millimeters of water per cubic meter of air)	4.8	6.8	9.4	17.3	23.1	30.4

- a. Find an exponential regression equation for these data. Round to the nearest thousandth.  
 b. Use the equation to predict the number of milliliters of water per cubic meter of air at a temperature of 15°C. Round to the nearest tenth.
38. Artificial snow is made at a ski resort by combining air and water in a ratio that depends on the outside air temperature. The table below shows the rate of air flow needed for various temperatures.

Temperature (in °F)	0	5	10	15	20
Air Flow (in cubic feet per minute)	3.0	3.6	4.7	6.1	9.9

- a. Find an exponential regression equation for these data. Round to the nearest hundredth.  
 b. Use the equation to predict the air flow needed when the temperature is 25°F. Round to the nearest tenth.

### Extensions

#### CRITICAL THINKING

An exponential model for population growth or decay can be accurate over a short period of time. However, this model begins to fail because it does not account for the natural resources necessary to support growth, nor does it account for death within the population. Another model, called the *logistic model*, can account for some of these effects. The logistic model is given by

$$P(t) = \frac{mP_0}{P_0 + (m - P_0)e^{-kt}}$$

where  $P(t)$  is the population at time  $t$ ,  $m$  is the maximum population that can be sup-

ported,  $P_0$  is the population when  $t = 0$ , and  $k$  is a positive constant that is related to the growth of the population.

39. One model of Earth's population is given by  $P(t) = \frac{280}{4 + 66e^{-0.013t}}$ . In this equation,  $P(t)$  is the population in billions and  $t$  is the number of years after 1980. Round answers to the nearest hundred million.
- a. According to this model, what was Earth's population in the year 2000?  
 b. According to this model, what will be Earth's population in the year 2010?  
 c. If  $t$  is very large, say greater than 500, then  $e^{-0.013t} \approx 0$ . What does this suggest about the maximum population that Earth can support?
40. Biologists have determined that the maximum wolf population in a certain preserve is 1000 wolves. Suppose the population of wolves in the preserve in the year 2000 was 500, and that  $k$  is estimated to be 0.025.
- a. Find a logistic function for the number of wolves in the preserve in year  $t$ , where  $t$  is the number of years after 2000.  
 b. Find the estimated wolf population in 2015.

#### EXPLORATIONS

41. The formula used to calculate a monthly lease payment or a monthly car payment (for a purchase rather than a lease) is given by  $P = \frac{Ar(1+r)^n - Vr}{(1+r)^n - 1}$ , where  $P$  is the monthly payment,  $A$  is the amount of the loan,  $r$  is the monthly interest rate as a decimal,  $n$  is the number of months of the loan or lease, and  $V$  is the residual value of the car at the end of the lease. For a car purchase,  $V = 0$ .
- a. If the annual interest rate for a loan is 9%, what is the monthly interest rate as a decimal?  
 b. Write the formula for a monthly car payment when the car is purchased rather than leased.  
 c. Suppose you lease a car for 5 years. Find the monthly lease payment if the lease amount is \$10,000, the residual value is \$6000, and the annual interest rate is 6%.  
 d. Suppose you purchase a car and secure a 5-year loan for \$10,000 at an annual interest rate of 6%. Find the monthly payment.  
 e. Why are the answers to parts c and d different?




43. An airplane left Los Angeles at 8:20 A.M. and flew to Boston. The flying time was 6 hours 20 minutes. Boston is on Eastern Standard Time (EST) and Los Angeles is on Pacific Standard Time (PST), which is 3 hours behind EST. After the plane was on the ground for 1 hour it flew to Chicago, which is on Central Standard Time (CST). CST is 1 hour behind EST. The flying time from Boston to Chicago was 2 hours 20 minutes.
- a. What time, EST, did the plane arrive in Boston?  
 b. What time, CST, did the plane arrive in Chicago?
44. a. List the four steps in Polya's problem-solving strategy.  
 b. List eight problem-solving procedures that one might use in Polya's second step.

### Extensions

#### CRITICAL THINKING

45. What is the 100th decimal digit in the decimal representation of  $\frac{2}{3}$ ?
46. a. How many times larger is  $3^{(3^3)}$  than  $(3^3)^3$ ?  
 b. How many times larger is  $4^{(4^4)}$  than  $(4^4)^4$ ? *Note:* Most calculators will not display the answer to this problem because it is too large. However, the answer can be determined in exponential form by applying the following properties of exponents.
- $$(a^m)^n = a^{mn} \quad \text{and} \quad \frac{a^m}{a^n} = a^{m-n}$$
47. The mathematician Augustus De Morgan once wrote that he had the distinction of being  $x$  years old in the year  $x^2$ . He was 43 in the year 1849.
- a. Explain why people born in the year 1980 might share the distinction of being  $x$  years old in the year  $x^2$ . *Note:* Assume  $x$  is a natural number.  
 b. What is the next year after 1980 for which people born in that year might be  $x$  years old in the year  $x^2$ ?

48. Select a two-digit number between 50 and 100. Add 83 to your number. From this number form a new number by adding the digit in the hundreds place to the number formed by the other two digits (the digits in the tens place and the ones place). Now subtract this newly formed number from your original number. Your final result is 16. Use a deductive approach to show that the final result is always 16 regardless of which number you start with.
49. How many digits does it take in total to number a book from page 1 to page 240?
50. Consider a checkerboard with two red squares on opposite corners removed, as shown in the accompanying figure. Determine whether it is possible to completely cover the checkerboard with 31 dominos if each domino is placed horizontally or vertically and each domino covers exactly two squares. If it is possible, show how to do it. If it is not possible, explain why it cannot be done.



#### COOPERATIVE LEARNING

51. The object of this exercise is to create mathematical expressions that use exactly four 4's and that simplify to a counting number from 1 to 20, inclusive. You are allowed to use the following mathematical symbols:  $+$ ,  $-$ ,  $\times$ ,  $\div$ ,  $\sqrt{\quad}$ ,  $(\quad)$ , and  $!$ . For example,
- $$\frac{4}{4} + \frac{4}{4} = 2, \quad 4^{(4-4)} + 4 = 5, \quad \text{and}$$
- $$4 - \sqrt{4} + 4 \times 4 = 18$$
52. The following puzzle is a famous cryptarithm.

**SEND  
+ MORE  
= MONEY**

Each letter in the cryptarithm represents one of the digits 0 through 9. The leading digits, represented by

## Extensions

Extension exercises are places at the end of each exercise set. In most cases these exercises are more challenging and require more time and effort than the preceding exercises. The Extension exercises always include at least two of the following types of exercises:

- Critical Thinking*
- Cooperative Learning*
- Explorations*

Some Critical Thinking exercises require the application of two or more procedures or concepts.

The Cooperative Learning exercises are designed for small groups of 2–4 students.

Many of the Exploration exercises require students to search on the Internet or through reference materials in a library.

## Chapter Summary

At the end of each chapter there is a Chapter Summary that includes Key Terms and Essential Concepts that were covered in the chapter. These chapter summaries provide a single point of reference as the student prepares for an examination. Each key word references the page number of the chapter where the word was first introduced.

### CHAPTER 1 Summary

#### Key Terms

conjecture [p. 4]  
 counterexample [p. 7]  
 difference table [p. 15]  
 Fibonacci sequence [p. 19]  
 first, second, and third differences [p. 15]  
 $n$ th term formula [p. 17]  
 $n$ th term of a sequence [p. 15]  
 palindromic number [p. 39]  
 Pascal's Triangle [p. 38]  
 prime number [p. 13]  
 recursive definition [p. 20]  
 sequence [p. 14]  
 term of a sequence [p. 14]

#### Essential Concepts

Inductive reasoning is the process of reaching a general conclusion by examining specific examples. A conclusion based on inductive reasoning is called a *conjecture*. A conjecture may or may not be correct.

- Deductive reasoning is the process of reaching a conclusion by applying general assumptions, procedures, or principles.
- A statement is a *true statement* provided it is true in all cases. If you can find one case in which a statement is not true, called a *counterexample*, then the statement is a *false statement*.
- The terms of the *Fibonacci sequence* 1, 1, 2, 3, 5, 8, 13, 21, ... can be determined by using the *recursive definition*  
 $F_1 = 1, F_2 = 1, \text{ and } F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 3$
- Many problems can be solved by applying *Polya's problem solving strategy*:
  - Understand the problem.
  - Devise a plan.
  - Carry out the plan.
  - Review your solution.

page 42


### Chapter 1 • Review Exercises 43

#### CHAPTER 1 Review Exercises

In Exercises 1–4, determine whether the argument is an example of inductive reasoning or deductive reasoning.

- All books written by J. K. Rowling make the best-seller list. The book *Harry Potter and the Goblet of Fire* is a J. K. Rowling book. Therefore, *Harry Potter and the Goblet of Fire* made the best-seller list.
- Samantha got an A on each of her first four math tests, so she will get an A on the next math test.
- We had rain yesterday, so there is less chance of rain today.
- All amoeba multiply by dividing. I have named the amoeba shown in my microscope Amelia. Therefore, Amelia multiplies by dividing.
- Find a counterexample to show that the following conjecture is false.  
 Conjecture: For all  $x$ ,  $x^4 > x$ .
- Find a counterexample to show that the following conjecture is false.  
 Conjecture: For all counting numbers  $n$ ,  $\frac{n^2 + 5n + 6}{n}$

In Exercises 13 and 14, determine the  $n$ th term formula for the number of square tiles in the  $n$ th figure.



page 43

## Chapter Review




Review exercises are found near the end of each chapter. These exercises were selected to help the student integrate the major topics presented in the chapter. The answers to all of the Chapter Review exercises appear in the answer section along with a section reference for each exercise. These section references indicate the section or sections where a student can locate the concepts needed to solve each exercise.

## Chapter Test

The Chapter Test exercises are designed to simulate a possible test of the material in the chapter. The answers to all of the Chapter Test exercises appear in the answer section along with a section reference for each exercise. The section references indicate the section or sections where a student can locate the concepts needed to solve each exercise.

### CHAPTER 8 Test

- Find the volume of a cylinder with a height of 6 m and a radius of 3 m. Round to the nearest hundredth.
- Find the perimeter of a rectangle that has a length of 2 m and a width of 1.4 m.
- Find the complement of a  $32^\circ$  angle.
- Find the area of a circle that has a diameter of 2 m. Round to the nearest hundredth.
- In the figure below, lines  $l_1$  and  $l_2$  are parallel. Angle  $x$  measures  $30^\circ$ . Find the measure of angle  $y$ .
- In the figure below, lines  $l_1$  and  $l_2$  are parallel. Angle  $x$  measures  $45^\circ$ . Find the measures of angles  $a$  and  $b$ .
- Find the area of a square that measures 2.25 ft on each side.
- Find the volume of the figure.



page 543